

REVIEW ON NEW PHYSICS IN HEAVY FLAVORS: B -MESON SECTOR

The off-diagonal elements M_{12}^s and Γ_{12}^s can be related to the physical observables in the B_s -meson sector by introducing the following model-independent parameterization:

$$\begin{aligned} M_{12}^s &= (M_{12}^s)_{\text{SM}} + (M_{12}^s)_{\text{NP}} = (M_{12}^s)_{\text{SM}} R_M e^{i\phi_M}, \\ \Gamma_{12}^s &= (\Gamma_{12}^s)_{\text{SM}} + (\Gamma_{12}^s)_{\text{NP}} = (\Gamma_{12}^s)_{\text{SM}} R_\Gamma e^{i\phi_\Gamma}. \end{aligned} \quad (2)$$

To leading power in $|\Gamma_{12}^s|/|M_{12}^s|$ the mass difference ΔM_s , the CP-violating phase $\phi_{J/\psi\phi}^s$, the decay-width difference $\Delta\Gamma_s$, and the flavor-specific CP asymmetry a_{fs}^s are then given by

$$\begin{aligned} \Delta M_s &= (\Delta M_s)_{\text{SM}} R_M, \quad \phi_{J/\psi\phi}^s = (\phi_{J/\psi\phi}^s)_{\text{SM}} + \phi_M, \\ \Delta\Gamma_s &\approx (\Delta\Gamma_s)_{\text{SM}} R_\Gamma \cos(\phi_M - \phi_\Gamma), \quad a_{fs}^s \approx (a_{fs}^s)_{\text{SM}} \frac{R_\Gamma}{R_M} \frac{\sin(\phi_M - \phi_\Gamma)}{\phi_{\text{SM}}^s}, \end{aligned} \quad (3)$$

with $\phi_{\text{SM}}^s = (0.22 \pm 0.06)^\circ$.⁴

The Λ_{NP} -scaling (1) combined with (2) and (3) suggests that new particles are more likely to leave a visible imprint in ΔM_s and $\phi_{J/\psi\phi}^s$ than in $\Delta\Gamma_s$ and a_{fs}^s . Experimentally, however, the mass difference and the CP phase in B_s - $\bar{B}_s$ mixing are spot on the SM predictions,¹ while a_{fs}^s , if extracted from the latest measurement⁵ of the like-sign dimuon charge asymmetry A_{SL}^b , assuming the absence of NP in the B_d -meson system, shows an anomaly of almost 4σ . In view of these results, it seems worthwhile to ask: how big can NP in Γ_{12}^s be afterall?

While any operator $(\bar{s}b)f$ with f leading to a flavor-neutral final state of two or more fields and mass below the bottom-quark threshold can alter Γ_{12}^s , the possible final states are in practice limited, because essentially all $B_s \rightarrow f$ and $B_d \rightarrow X_s f$ decay modes involving light states in the final state are strongly constrained experimentally. An exception are B decays to tau pairs.⁶ The possibility of large $b \rightarrow s\tau^+\tau^-$ contributions to Γ_{12}^s , can be analyzed in a model-independent fashion⁷ by adding the complete set of dimension-six operators ($A, B = L, R$)

$$\begin{aligned} Q_{S,AB} &= (\bar{s} P_A b) (\bar{\tau} P_B \tau), \\ Q_{V,AB} &= (\bar{s} \gamma^\mu P_A b) (\bar{\tau} \gamma_\mu P_B \tau), \\ Q_{T,A} &= (\bar{s} \sigma^{\mu\nu} P_A b) (\bar{\tau} \sigma_{\mu\nu} P_A \tau), \end{aligned} \quad (4)$$

to the SM Lagrangian. Here $P_{L,R} = (1 \mp \gamma_5)/2$ project onto left- and right-handed chiral fields and $\sigma^{\mu\nu} = i[\gamma^\mu, \gamma^\nu]/2$.

The ten operators entering (4) govern the purely leptonic $B_s \rightarrow \tau^+\tau^-$ decay, the inclusive semileptonic $B \rightarrow X_s \tau^+\tau^-$ decay, and its exclusive counterpart $B^+ \rightarrow K^+ \tau^+\tau^-$, making these channels potentially powerful probes of $(\bar{s}b)(\bar{\tau}\tau)$ operators. In practice, these direct constraints however turn out to be rather loose at present. Explicitly, one obtains

$$\begin{aligned} \text{BR}(B_s \rightarrow \tau^+\tau^-) &< 3\%, \\ \text{BR}(B \rightarrow X_s \tau^+\tau^-) &\lesssim 2.5\%, \\ \text{BR}(B^+ \rightarrow K^+ \tau^+\tau^-) &< 3.3 \cdot 10^{-3}. \end{aligned} \quad (5)$$

Here the first limit derives⁸ from comparing the SM prediction $\tau_{B_s}/\tau_{B_d} - 1 \in [-0.4, 0.0]\%$ ⁹ with the corresponding experimental result $\tau_{B_s}/\tau_{B_d} - 1 = (0.4 \pm 1.9)\%$, while the second (crude) bound follows from estimating⁷ the possible contamination of the exclusive and inclusive semileptonic decay samples by $B \rightarrow X_s \tau^+\tau^-$ events. The final number corresponds to the 90% confidence level (CL) upper limit on the branching ratio of $B^+ \rightarrow K^+ \tau^+\tau^-$ as measured by BaBar.¹⁰

Further constraints on the Wilson coefficients of the $(\bar{s}b)(\bar{\tau}\tau)$ operators arise indirectly from the experimentally available information on the $b \rightarrow s\gamma$, $b \rightarrow s\ell^+\ell^-$ ($\ell = e, \mu$), and $b \rightarrow s\gamma\gamma$ transitions, because some of the effective operators introduced in (4) mix into the electromagnetic

dipole operators $Q_{7,A}$ and the vector-like semileptonic operators $Q_{9,A}$. An explicit calculation⁷ shows that the operators $Q_{S,AB}$ mix neither into $Q_{7,A}$ nor $Q_{9,A}$, while $Q_{V,AB}$ ($Q_{T,A}$) mixes only into $Q_{9,A}$ ($Q_{7,A}$). As a result of the particular mixing pattern, the stringent constraints from the radiative decay $B \rightarrow X_s \gamma$ rule out large contributions to Γ_{12}^s only if they arise from the tensor operators $Q_{T,A}$. Similarly, the rare decays $B \rightarrow X_s \ell^+ \ell^-$ and $B \rightarrow K^{(*)} \ell^+ \ell^-$ primarily limit contributions stemming from the vector operators $Q_{V,AB}$. In contrast to $B \rightarrow X_s \gamma$, all $(\bar{s}b)(\bar{\tau}\tau)$ operators contribute to the double-radiative $B_s \rightarrow \gamma\gamma$ decay at the one-loop level. A detailed study⁷ shows however that the limits following from $b \rightarrow s\gamma\gamma$ are in practice not competitive with the bounds obtained from the other tree- and loop-level mediated $B_{s,d}$ -meson decays.

The off-diagonal element Γ_{12}^s is related via the optical theorem to the absorptive part of the forward-scattering amplitude which converts a $\bar{B}_s$ into a B_s meson. Working to leading order in the strong coupling constant and Λ_{QCD}/m_b , the contributions from the operators (4) to Γ_{12}^s is found by computing the matrix elements of the (Q_i, Q_j) double insertions between quark states. Such a calculation⁷ leads to the following 90% CL bounds

$$(R_\Gamma)_{S,AB} < 1.15, \quad (R_\Gamma)_{V,AB} < 1.35, \quad (R_\Gamma)_{T,L} < 1.004, \quad (R_\Gamma)_{T,R} < 1.008. \quad (6)$$

These numbers imply that $(\bar{s}b)(\bar{\tau}\tau)$ operators of scalar (vector) type can lead to enhancements of $|\Gamma_{12}^s|$ over its SM value by 15% (35%) without violating any existing constraint. In contrast, contributions from tensor operators can alter $|\Gamma_{12}^s|$ by at most 0.8%. Since an explanation of the experimentally observed large negative values of a_{fs}^s (or equivalent A_{SL}^b) calls for NP in Γ_{12}^s that changes the SM value by a factor of 3 or more, absorptive NP in the form of $(\bar{s}b)(\bar{\tau}\tau)$ operators obviously does not provide a satisfactory description of the large dimuon charge asymmetry observed by the DØ collaboration. This is a model-independent conclusion that can be shown to hold in explicit models of NP with modification of the $b \rightarrow s\tau^+\tau^-$ channel such as leptoquark scenarios or Z' models.⁷

3 Interplay between high- and low- p_T measurements

The discovery of the Higgs boson with mass of around 125 GeV^{11,12} combined with the direct limits on new particles rule out most of the simple and natural implementations of NP. At this stage, it is hence very important to examine the plethora of available experimental data and to look for hints that can guide us towards special regions where NP may still hide. Indirect probes of NP as provided by rare B -meson decays can play a key role in this endeavor, as will be described in the following for the case of the minimal supersymmetric SM (MSSM).

In the decoupling limit of the MSSM, *i.e.*, $M_A^2 \gg M_Z^2$, the lightest CP-even Higgs boson acquires the loop-corrected squared mass¹³

$$M_h^2 \approx M_Z^2 c_{2\beta}^2 + \frac{3G_F}{\sqrt{2}\pi^2} m_t^4 \left[-\ln\left(\frac{m_t^2}{m_{\tilde{t}}^2}\right) + \frac{X_t^2}{m_{\tilde{t}}^2} \left(1 - \frac{X_t^2}{12m_{\tilde{t}}^2}\right) \right]. \quad (7)$$

Here $c_{2\beta} = \cos(2\beta)$, $m_{\tilde{t}}^2 = m_{\tilde{t}_1} m_{\tilde{t}_2}$, and $X_t = A_t - \mu/t_\beta$ denotes the stop-mixing parameter, which depends on the trilinear stop-Higgs boson coupling A_t and the higgsino mass parameter μ . From the above expression, one concludes that to raise M_h from the Z -boson mass M_Z to 125 GeV requires sizable values of $t_\beta = \tan\beta$, a heavy stop spectrum ($m_{\tilde{t}} \gtrsim 1$ TeV), and/or large stop mixing ($|A_t| \gtrsim 2$ TeV).

The mentioned MSSM parameters also play an important role in the production and the decay of the Higgs. Applying Higgs low-energy theorems, it is easy to derive that the modification of the $gg \rightarrow h$ production cross section due to stops, can be written as¹⁴

$$R_h = \frac{\sigma(gg \rightarrow h)_{\text{MSSM}}}{\sigma(gg \rightarrow h)_{\text{SM}}} \approx \left[1 + \frac{m_{\tilde{t}}^2}{4} \left(\frac{1}{m_{\tilde{t}_1}^2} + \frac{1}{m_{\tilde{t}_2}^2} - \frac{X_t^2}{m_{\tilde{t}_1}^2 m_{\tilde{t}_2}^2} \right) \right]^2. \quad (8)$$




Figure 1: Predictions for M_h in GeV (upper left), R_γ (upper center), R_{X_s} (upper right), $R_{\mu+\mu^-}$ (lower left), Δa_μ (lower center), and Ωh^2 (lower right). The dotted black lines indicate the parameter regions with $R_\gamma > 1$, while the dashed black lines correspond to the 95% CL regions favored by $B \rightarrow X_s \gamma$. See text for further explanations.

One infers that the amount of mixing in the stop sector determines whether (8) is smaller or larger than 1. For no mixing ($X_t = 0$), one has $R_h > 1$, while if X_t is parametrically larger than the mass eigenvalues $m_{\tilde{t}_{1,2}}$ (with $m_{\tilde{t}_1} < m_{\tilde{t}_2}$) then one has $R_h < 1$. The fact that in the MSSM, to make the Higgs sufficiently heavy, one needs large/maximal mixing, then implies that for a random MSSM parameter point with $M_h \approx 125$ GeV one should find a suppression of $\sigma(gg \rightarrow h)$. In fact, this is precisely what happens in large parts of the parameter space.

For $M_A \gg M_Z$ and $t_\beta \gg 1$, charged Higgs as well as chargino effects are strongly suppressed in $h \rightarrow \gamma\gamma$, but stau loops can have a notable impact on the diphoton signal strength

$$R_\gamma = \frac{[\sigma(pp \rightarrow h)\text{BR}(h \rightarrow \gamma\gamma)]_{\text{MSSM}}}{[\sigma(pp \rightarrow h)\text{BR}(h \rightarrow \gamma\gamma)]_{\text{SM}}} \approx 1 + 0.10 \frac{m_\tau^2 X_\tau^2}{m_{\tilde{\tau}_1}^2 m_{\tilde{\tau}_2}^2}, \quad (9)$$

if mixing in the stau sector is large, *i.e.*, $|X_\tau| = |A_\tau - \mu t_\beta| \gg m_{\tilde{\tau}_{1,2}}$ and the stau mass eigenstate $\tilde{\tau}_1$ is sufficiently light.¹⁵ In (9) we have assumed that $M_h \approx 125$ GeV and explicitly included only stau effects.

The above discussion should have made clear that the part of the MSSM parameter space with $M_A, t_\beta \rightarrow \infty$ represents a phenomenologically interesting region for Higgs physics. How do the dominant MSSM corrections to $B \rightarrow X_s \gamma$ and $B_s \rightarrow \mu^+ \mu^-$ look like in this limiting case? For $B \rightarrow X_s \gamma$, one obtains¹⁶

$$R_{X_s} = \frac{\text{BR}(B \rightarrow X_s \gamma)_{\text{MSSM}}}{\text{BR}(B \rightarrow X_s \gamma)_{\text{SM}}} \approx 1 - 2.61 \Delta C_7 + 1.66 (\Delta C_7)^2, \quad (10)$$

where the dominant one-loop contributions are provided by diagrams with top squarks and higgsino-like charginos, leading to $\Delta C_7 \propto -\mu A_t t_\beta m_t^2/m_{\tilde{t}}^4$. The sign of ΔC_7 combined with that of (10) implies that for $\mu A_t > 0$ ($\mu A_t < 0$) the MSSM branching ratio is larger (smaller) than the SM expectation. For the choices $t_\beta = 50$, $m_{\tilde{t}} = 1.5$ TeV, $|\mu| = 1$ TeV, and $|A_t| = 3$ TeV,

one finds numerically an enhancement/suppression of $\text{BR}(B \rightarrow X_s \gamma)_{\text{MSSM}}$ by $\mathcal{O}(30\%)$. Shifts of this size are detectable given the present theoretical calculations and experimental extractions.

In the case of the purely leptonic B_s decay, one can write¹⁶

$$R_{\mu^+\mu^-} = \frac{\text{BR}(B_s \rightarrow \mu^+\mu^-)_{\text{MSSM}}}{\text{BR}(B_s \rightarrow \mu^+\mu^-)_{\text{SM}}} \approx 1 - 13.2 C_P + 43.6 (C_S^2 + C_P^2), \quad (11)$$

where $C_{P,S}$ denote the dimensionless Wilson coefficients of the semileptonic pseudo-scalar and scalar operators. The term linear in C_P arises from the interference with the SM contribution to the semileptonic axial-vector operator. For $C_P > 0$ it interferes destructively with the term proportional to $(C_S^2 + C_P^2)$, which implies that a pseudo-scalar contribution of the correct sign and size will lead to a suppression of $\text{BR}(B_s \rightarrow \mu^+\mu^-)_{\text{MSSM}}$ below its SM value.

Within the MSSM the contributions to $C_{P,S}$ with the strongest t_β dependence arise from neutral Higgs double penguins.¹⁷ In the decoupling limit, one has

$$C_P \approx -C_S \propto \mu A_t \frac{t_\beta^3}{(1 + \epsilon_b t_\beta)^2} \frac{m_t^2}{m_{\tilde{t}}^2} \frac{m_b m_\mu}{M_W^2 M_A^2}, \quad \epsilon_b \propto \frac{\alpha_s}{\pi} \frac{\mu M_3}{m_{\tilde{t}}^2}. \quad (12)$$

Here ϵ_b encodes loop-induced non-holomorphic terms due to gluino exchange which introduce a dependence of $C_{P,S}$ on $\text{sgn}(\mu M_3)$. From (12) one deduces that the sign of C_S (C_P) is opposite to (follows) that of μA_t and that both coefficients are suppressed (enhanced) for $\mu M_3 > 0$ ($\mu M_3 < 0$). The recent LHCb measurement³ of $\text{BR}(B_s \rightarrow \mu^+\mu^-) = (3.2_{-1.2}^{+1.5}) \cdot 10^{-9}$ hence clearly favors $\mu A_t > 0$ and $\mu M_3 > 0$, if Higgs exchange gives the dominant contribution to (11).

In Figure 1 we show the results of numerical scans in the A_t - μ plane.¹⁶ We restrict ourselves to the quadrant with $A_t > 0$ and $\mu > 0$ since it shows the most interesting effects and correlations. The predictions correspond to $t_\beta = 60$, $M_A = 1 \text{ TeV}$, $M_1 = 50 \text{ GeV}$, $M_2 = 300 \text{ GeV}$, $M_3 = 1.2 \text{ TeV}$, $\tilde{m}_{Q_3} = \tilde{m}_{u_3} = 1.5 \text{ TeV}$, $\tilde{m}_{L_3} = \tilde{m}_{l_3} = 350 \text{ GeV}$, while we take common soft masses of 1.5 TeV and 2 TeV (1 TeV) for the other “left-” and “right-handed” squark (sleptons). We furthermore employ $A_b = 2.5 \text{ TeV}$ and $A_\tau = 500 \text{ GeV}$, while the first and second generation trilinear couplings take the same values as those of the third generation. We see that for the above choice of parameters, A_t has to lie in the range of $[2, 5] \text{ TeV}$ to accommodate $M_h \in [123, 129] \text{ GeV}$ and that $R_\gamma > 1$ can only be obtained in a narrow sliver around $\mu = 900 \text{ GeV}$. For our choice of soft masses $\tilde{m}_{L_3}$ and $\tilde{m}_{l_3}$ such large μ values lead to stau masses $m_{\tilde{\tau}_1} \in [80, 120] \text{ GeV}$. This strong correlation between notable enhancements in R_γ and the presence of a very light stau is a smoking gun signal of the discussed scenario. From the panel showing R_{X_s} , one infers that $\text{BR}(B \rightarrow X_s \gamma)$ is always enhanced (by about 20% to 60%) with respect to the SM. This is an interesting and potentially important finding, since the $B \rightarrow X_s \gamma$ constraint starts cutting into the already narrow region in the A_t - μ plane with $M_h \approx 125 \text{ GeV}$ and $R_\gamma > 1$. We also observe that solutions that satisfy $B \rightarrow X_s \gamma$ typically feature a suppression of $B_s \rightarrow \mu^+\mu^-$. In fact, asking for an agreement with R_{X_s} at the 95% CL as well as $R_\gamma > 1$, implies $R_{\mu^+\mu^-} \in [0.6, 1.0]$. The figure furthermore shows that assuming a light slepton spectrum, the long-standing discrepancy of the anomalous magnetic moment of the muon a_μ is reduced in the parameter region selected by M_h and the enhanced diphoton signal. Finally, notice that in this very region of parameter space also the correct thermal dark matter relic density Ωh^2 can be achieved, but only if one assumes the hierarchy $|M_1| \ll |M_2| \ll |\mu|$. A typical MSSM spectrum leading to a significantly enhanced $h \rightarrow \gamma\gamma$ rate as well as the correct value of Ωh^2 , hence contains a light bino as the dark matter candidate, a light and maximally mixed stau, and a heavy higgsino. The aforementioned deviations and found correlations could be tested in the near future and hence may become very valuable as guidelines and consistency checks, in particular if the high- p_T LHC experiments start to see supersymmetric partners.

4 Conclusions

While the 7 TeV and 8 TeV LHC runs took the hope to find spectacular effects in heavy flavor physics, the low- p_T data is at present not precise enough to exclude NP contaminations of $\mathcal{O}(50\%)$. This still leaves room for visible and interesting effects in rare B -meson decays, given the theoretical cleanness of these observables. Since there is no direct sign of NP at the LHC and also the Higgs signal strengths look very much SM-like, such indirect NP probes are more important than ever and only the synergy between high- and low- p_T observations may give us the key to solving the puzzles of fundamental physics. The expected LHC precision measurements of the B -mixing observables, $B_s \rightarrow \mu^+ \mu^-$, $B \rightarrow K^* \ell^+ \ell^-$, the angle γ , *etc.* may play a crucial role in this endeavor.

Acknowledgments

I am grateful to the organizers of the BSM conference Vietnam 2012 for the invitation to a great conference, and to Marc Besancon for encouraging me to write these proceedings. Travel support from the UNILHC network (PITN-GA-2009-237920) is acknowledged.

References

1. P. Uwer, Talk given at La Thuile 2013, <https://cds.cern.ch/record/1519663/files/LHCb-TALK-2013-040.pdf>
2. H. Ruiz, Talk given at La Thuile 2013, <https://agenda.infn.it/getFile.py/access?contribId=27&tsessionId=8&resId=0&materialId=slides&confId=5540>
3. R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **110**, 021801 (2013) [arXiv:1211.2674 [hep-ex]].
4. A. Lenz *et al.*, Phys. Rev. D **83**, 036004 (2011) [arXiv:1008.1593 [hep-ph]].
5. V. M. Abazov *et al.* [DØ Collaboration], Phys. Rev. D **84**, 052007 (2011) [arXiv:1106.6308 [hep-ex]].
6. A. Dighe, A. Kundu and S. Nandi, Phys. Rev. D **76**, 054005 (2007) [arXiv:0705.4547 [hep-ph]].
7. C. Bobeth and U. Haisch, arXiv:1109.1826 [hep-ph].
8. A. Dighe, A. Kundu and S. Nandi, Phys. Rev. D **82**, 031502 (2010) [arXiv:1005.4051 [hep-ph]].
9. A. Lenz, arXiv:1205.1444 [hep-ph] and references therein.
10. K. Flood [BaBar Collaboration], PoS **ICHEP2010**, 234 (2010).
11. G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012) [arXiv:1207.7214 [hep-ex]].
12. S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012) [arXiv:1207.7235 [hep-ex]].
13. A. Djouadi, Phys. Rept. **459**, 1 (2008) [arXiv:hep-ph/0503173] and references therein.
14. R. Dermisek and I. Low, Phys. Rev. D **77**, 035012 (2008) [arXiv:hep-ph/0701235].
15. M. Carena, S. Gori, N. R. Shah and C. E. M. Wagner, JHEP **1203**, 014 (2012) [arXiv:1112.3336 [hep-ph]].
16. U. Haisch and F. Mahmoudi, JHEP **1301**, 061 (2013) [arXiv:1210.7806 [hep-ph]].
17. K. S. Babu and C. F. Kolda, Phys. Rev. Lett. **84**, 228 (2000) [arXiv:hep-ph/9909476].